


## Low Temperatures Enhance Ozone Degradation above the Arctic

KIT Climate Researchers Studied Ozone Hole Formation - Results in "Geophysical Research Letters"

MIPAS O3 20110318 50.00 hPa

MIPAS CLO am 20110318 50.00 hPa



March 2011: Strongly reduced ozone values (left, dark blue) and significantly increased concentration of chlorine monoxide (right, red) that is directly involved in ozone degradation. (Figure: IMK-ASF, KIT)

**Extraordinarily cold temperatures in the winter of 2010/2011 caused the most massive destruction of the ozone layer above the Arctic so far: The mechanisms leading to the first ozone hole above the North Pole were studied by scientists of the KIT Institute of Meteorology and Climate Research (IMK). According to these studies, further cooling of the ozone layer may enhance the influence of ozone-destroying substances, e.g. chlorofluorocarbons (CFC), such that repeated occurrence of an ozone hole above the Arctic has to be expected.**

About a year ago, IMK scientists, together with colleagues from Oxford, detected that ozone degradation above the Arctic for the first time reached an extent comparable to that of the ozone hole above the South Pole. Then, the KIT researchers studied the mechanisms behind. Their results have now been published in the journal "Geo-

**Monika Landgraf**  
Press Officer

Kaiserstraße 12  
76131 Karlsruhe, Germany  
Phone: +49 721 608-4 7414  
Fax: +49 721 608-4 3658

**For further information,  
please contact:**

Margarete Lehné  
Public Relations and  
Marketing (PKM)  
Phone: +49 721 608-4 8121  
Fax: +49 721 608-4 3658  
E-mail: margarete.lehne@kit.edu

physical Research Letters”.

According to IMK studies, occurrence of the Arctic ozone hole was mainly due to the extraordinarily cold temperatures in the ozone layer that is located at about 18 km height in the stratosphere, i.e. the second layer of the earth’s atmosphere. There, chlorine compounds originating from chlorofluorocarbons (CFC, e.g. greenhouse gases and refrigerants) and other pollutants are converted chemically at temperatures below  $-78^{\circ}\text{C}$ . These chemical conversion products attack the ozone layer and destroy it partly. One of the main statements in the study: If the trend to colder temperatures in the stratosphere observed in the past decades will continue, repeated occurrence of an Arctic ozone hole has to be expected.

The team of IMK researchers analyzed measurements of the chemical composition of the atmosphere by the MIPAS satellite instrument developed by KIT. In addition, model calculations were made to determine concrete effects of further cooling of the ozone layer. “We found that further decrease in temperature by just  $1^{\circ}\text{C}$  would be sufficient to cause a nearly complete destruction of the Arctic ozone layer in certain areas,” says Dr. Björn-Martin Sinnhuber, main author of the study. Observations over the past thirty years indicate that the stratosphere in cold Arctic winters cooled down by about  $1^{\circ}\text{C}$  per decade on the average. According to Sinnhuber, further development of the ozone layer will consequently be influenced also by climate change. He points out that the increase in carbon dioxide and other greenhouse gases will warm up the bottom air layers near the ground due to the reflection of part of the thermal radiation by the bottom layer of the atmosphere towards the earth’s surface, but also result in a cooling of the air layers of the stratosphere above, where the ozone layer is located.

After the first discovery of the Antarctic ozone hole in the mid-1980s, CFCs were rapidly identified to be the cause and their use was prohibited by the Montreal Protocol of 1987. However, it will take decades until these substances will have been removed completely from the atmosphere. “Future cooling of the stratosphere would enhance and extend the impacts of these substances on the ozone layer,” says Dr. Björn-Martin Sinnhuber. It is now necessary to study potential feedbacks on climate change.

The present study is embedded in long-term programs of IMK in this field. In December, the researchers started a new measurement campaign in the Arctic ozone layer in Northern Sweden using a

high-altitude aircraft. Again, they encountered extraordinarily low temperatures. However, it is not yet possible to predict whether temperatures will be low enough over a longer term to cause a comparably large degradation of ozone in this winter.

**More information:** Geophysical Research Letters, volume 38, doi:10.1029/2011GL049784.

**Karlsruhe Institute of Technology (KIT) is a public corporation according to the legislation of the state of Baden-Württemberg. It fulfills the mission of a university and the mission of a national research center of the Helmholtz Association. KIT focuses on a knowledge triangle that links the tasks of research, teaching, and innovation.**

This press release is available on the internet at [www.kit.edu](http://www.kit.edu).

The photo of printing quality may be downloaded under [www.kit.edu](http://www.kit.edu) or requested by mail to [presse@kit.edu](mailto:presse@kit.edu) or phone +49 721 608-4 7414. The photo may be used in the context given above exclusively.